


Flow-Ecology Relationships

Upper Savannah RBC, Nov 8th, 2023




Drs. Luke Bower, Joe Mruzek, and Brandon Peoples

Using aquatic organisms to learn about river health



Too much water to monitor!



- >28,000 segments in SC
- >15,000 river miles
- And that's just wadeable streams (~84% of surface water in SC)


Too much water to monitor!



for people to



SC Freshwater Diversity



- 146* fish species
- 1,092 invertebrate groups (many more species)

Bio-assessment: using aquatic organisms to learn about river health

ASSESSMENT OF BIOTIC INTEGRITY USING FISH COMMUNITIES

James R. Karr

ABSTRACT

Man's activities have had profound, and usually negative, influences on freshwater fishes from the smallest streams to the largest rivers. Some negative effects are due to contaminants, while others are associated with changes in watershed hydrology, habitat modifications, and alteration of energy sources upon which the aquatic biota depends. Regrettably, past efforts to evaluate effects of man's activities on fishes have attempted to use water quality as a surrogate for more comprehensive biotic assessment. A more refined biotic assessment program is required for effective protection of freshwater fish resources. An assessment system proposed here uses a series of fish community attributes related to species composition and ecological structure to evaluate the quality of an aquatic biota. In preliminary trials this system accurately reflected the status of fish communities and the environment supporting them.

Passage of the Water Quality Act Amendments of 1972 (PL 92-500) stimulated many efforts to monitor the quality of water resource systems. Unfortunately, these efforts concentrated on development of thresholds and criteria levels for specific contaminants, often based on acute toxicity tests. The use of these criteria has been attacked on numerous grounds (Thurston et al. 1979); for example, they have not taken into account naturally occurring geographic variation of contaminants (e.g., asbestos, iron, zinc), considered the synergistic effects of numerous contaminants, nor considered sublethal effects (e.g., reproduction, growth) of most contaminants. In addition, monitoring of water quality parameters (nutrients, DO, temperature, pesticides, heavy metals, and other toxics) often misses short-term events that may be critical to assessment of biotic impacts. Finally, it is impossible to measure all factors that may impact biotic integrity. In fact, much literature on chem-



James R. Karr

THE AUTHOR: James R. Karr is Professor of Ecology in the Department of Ecology, Ethology, and Evolution and the Department of Forestry at the University of Illinois at Urbana-Champaign, and an affiliate of the Aquatic Biology Section of the Illinois Natural History Survey. He received a B.S. from Iowa State University and M.S. and Ph.D. from the University of Illinois. He has served as a consultant to USEPA, USDA-SCS, and Organization of American States (in Venezuela) on water resource and watershed development projects. His interest in fisheries develops from an interest in applying ecological principles to a variety of resource problems. Another major research focus deals with tropical forest ecology, especially bird communities. Address: Department of Ecology, Ethology, and Evolution, University of Illinois, 606 E. Healey, Champaign, Illinois 61820.

ical contaminants is of questionable value for setting water quality standards for aquatic organisms (Gosz 1980). Chemical monitoring misses many of the man-induced perturbations that impair use. For example, flow alterations, habitat degradation, heated effluents, and uses for power generation are not detected in chemical sampling. In short, criteria that emphasize physical and chemical attributes of water are unsuccessful as surrogates for measuring biotic integrity (Karr and Dudley 1981).


Recent legislation (Clean Water Act of 1977, PL 95-217) clearly calls for a more refined approach when pollution is defined as "the manmade or man-induced alteration of the chemical, physical, biological, and radiological integrity of water." Despite this refinement, regulatory agencies have been slow to replace the classical approach (uniform standards focusing on contaminant levels) with a more sophisticated and environmentally sound approach.

The integrity of water resources can best be assessed by evaluating the degree to which waters provide for beneficial uses. Important uses as defined by society may include water supply, recreational, and other uses as well as the preservation of future options for the use of the resource. Since an ability to sustain a balanced biotic community is one of the best indicators of the potential for beneficial use, sophisticated monitoring programs should seek to assess "biotic integrity."


This paper describes a procedure for monitoring water resources using fish. My contention is that by carefully monitoring fishes, one can rapidly assess the health ("biotic integrity") of a local water resource. In short, carefully planned monitoring and assessment can rapidly and relatively inexpensively serve as an exploratory assessment of water resource quality. Where impaired use is suggested by biological monitoring, a more nearly complete monitoring program can be implemented in search of the causative agent(s).

WHY MONITOR FISH?

Biological communities reflect watershed conditions since they are sensitive to changes in a wide array of environmental factors. Many groups of organisms have been proposed as indicators of environmental quality, but no single group has emerged as the






Flow-ecology relationships



Use of the relationships




-  Middle Atlantic Coastal Plain
-  Piedmont











Purpose


- To provide insight on the potential response of organisms to the alternate water withdrawal scenarios produced by SWAM.
 - We aim to put the SWAM results into a biological context.
- High demand water use scenario: 100 to 60 cfs



How will this work? Step 1




How will this work? Step 2



Ask RBC for a vote

How will this work? Step 3

Selected relationships




View SWAM results in a biological context

Scenario	Loss of species	Risk
MD	15%	Med
HD	25%	High

SWAM results

Scenario	Current	Predicted	% Change
MD	100	80	20%
HD	100	60	40%




Step 1: Quantify the flow-ecology relationships



Contents lists available at [ScienceDirect](#)

Science of the Total Environment

journal homepage: www.elsevier.com/locate/scitotenv




Quantifying flow–ecology relationships across flow regime class and ecoregions in South Carolina



Luke M. Bower ^{a,*}, Brandon K. Peoples ^b, Michele C. Eddy ^c, Mark C. Scott ^d

Framework

- The ecological limits of hydrologic alteration (ELOHA).
Poff et al., 2010



Build a hydrologic foundation of streamflow and biological data

B. Classify natural river types

C. Model and select flow ecology relationships


Biological Data:

- 492 Fish sites (streams & rivers)
 - DNR
 - 8 biological response metrics

- 530 aquatic insect sites
 - DHEC
 - 6 biological response metrics

Legend


- Fish sites
- ▲ Macroinvertebrate sites
- ▭ HUC6
- ▭ HUC8
- Blue Ridge
- Southern Coastal Plain
- Southeastern Plain
- Middle Atlantic Coastal Plain
- Piedmont



Characterizing aquatic diversity



- **Species richness:** number of species
- **Shannon's Diversity:** Accounts for percentages

Tolerant species



Diverse biota = healthy ecosystem

Hydrologic data



Legend

- Fish sites
- ▲ Macroinvertebrate sites

- HUC6
- HUC8
- Blue Ridge
- Southern Coastal Plain
- Southeastern Plain
- Middle Atlantic Coastal Plain
- Piedmont

Build a hydrologic foundation of streamflow data



- WaterFALL model:
 - rainfall-runoff model 30-year period
 - Accounts for withdrawals, discharges, and reservoirs within the river network
- 24 hydrologic metrics
- Flow regime: **Timing, magnitude, frequency, rate of change, and duration**

Table 2. Model Geospatial Inputs

Data Set	Name	Resolution	Reference
Hydrology	Enhanced National Hydrography Dataset Version 2	2.1 km ² within study area	Moore and Dewald, 2016
Land Cover	2016 National Land Cover Dataset	30-m grid	Jin et al., 2019
Climate	PRISM 4km Daily Temperature and Precipitation 1988–2018	4-km grid	PRISM Climate Group, 2019
Soils	Soil Survey Geographic Database (SSURGO)	1:12,000 to 1:63,360	USDA-NRCS, 2014
Subsurface Parameters	National Weather Service (NWS) for applications of the Sacramento Soil Moisture Accounting Model (SAC-SMA)	Approximately 4.7-km grid	Zhang et al., 2011

Received: 8 March 2021 | Revised: 23 September 2021 | Accepted: 25 October 2021

DOI: 10.1002/eco.2387

RESEARCH ARTICLE

WILEY

Predictability of flow metrics calculated using a distributed hydrologic model across ecoregions and stream classes: Implications for developing flow–ecology relationships

Michele C. Eddy¹  | Benjamin Lord¹  | Danielle Perrot¹ | Luke M. Bower²  |
Brandon K. Peoples³ 

Relevance of flow regime components

- Magnitude: MA1 (mean daily flow) and ML17 (base flow)
 - Alteration of habitat
 - Reduced water quality and higher mortality
- Duration: DL16 (duration of low water)
 - Alteration of connectivity
 - Increased duration of low water conditions
- Timing: TL1 (timing of low flow)
 - Loss of access to habitats
 - Disruption of life-cycle cues (spawning migration) and decreases in recruitment
 - Invasion of exotics



Framework

- The ecological limits of hydrologic alteration (ELOHA).
Poff et al., 2010

A. Build a hydrologic foundation of streamflow and biological data



Classify natural river types

C. Model and select flow ecology relationships

2. Classify natural river types

A. Flow-ecology relationships may differ among stream classes

A. Ecoregion

B. Hydrologic class




Framework

- The ecological limits of hydrologic alteration (ELOHA).
Poff et al., 2010

A. Build a hydrologic foundation of streamflow and biological data


B. Classify natural river types




Model and select flow ecology relationships

Identify relationships: some are informative


24 Flow metrics
14 Biotic metrics






Identify relationships: some are not informative



Identify relationships: remove uninformative relationships




Filter: statistical modeling process




Results summary

- We found >180 informative relationships across SC
 - Predicting responses
 - Defining biological response limits
- Many of these differed among stream classes
- All components of the flow regime were important to aquatic organisms
 - magnitude, frequency, duration, timing, and rate
- Next steps:
 - Identify those relevant to the Saluda
 - Present these proposed relationships to the RBC


Scenario	Loss of species	Risk
MD	15%	Med
HD	25%	High



How will this work? Step 1




How will this work? Step 2



How will this work? Step 3

Selected relationships




View SWAM results in a biological context

Scenario	Loss of species	Risk
MD	15%	Med
HD	25%	High

SWAM results

Scenario	Current	Predicted	% Change
MD	100	80	20%
HD	100	60	40%





Questions?

Imbower@clemson.edu